

**Application  
for  
United States Letters Patent**

**To all whom it may concern:**

Be it known that TUO JIN, LI CHEN AND HUA ZHU

have invented certain new and useful improvements in

STABLE POLYMER AQUEOUS/AQUEOUS EMULSION SYSTEM  
AND USES THEREOF

of which the following is a full, clear and exact description.

**STABLE POLYMER AQUEOUS/  
AQUEOUS EMULSION SYSTEM AND USES THEREOF**

5 Throughout this application, various references are referred to. Disclosures of these publications in their entirety are hereby incorporated by reference into this application to more fully describe the state of the art to which this  
10 invention pertains.

**BACKGROUND OF THE INVENTION**

15 While more and more biological therapeutic agents have become available due to the advances in molecular biology, immunology and microbiology, pharmaceutical development (e.g. development of appropriate dosage forms) for delivery of these  
20 proteins is behind the state of art of biotechnology. The situation is attributed to the difficulties in formulating these agents which are less permeable to tissue membranes, highly degradable at the sites of administration and  
25 therapy, short shelf life, and structurally susceptible during conventional formulation processes. To develop commercially available and patient compliant dosage forms for these therapeutics, the above-mentioned issues must be  
30 addressed respectively.

The disclosed are a novel material system named polymer aqueous/aqueous emulsions and its pharmaceutical and biotechnological applications. To  
35 meet desired therapeutic purposes such as sustained release, targeting to therapeutic sites, extension of bio-activity, and reducing toxicity, many

chemical and biological therapeutics need to be microencapsulated<sup>[1,2]</sup>. Emulsification is a key step in microencapsulation during which active ingredients are incorporated into the dispersed phase. Conventional emulsions are made by dispersing a hydrophilic phase (dispersed phase) into a hydrophobic phase (continuous phase) or vice versa (W/O or O/W)<sup>[3]</sup>. For microencapsulation of protein therapeutics, a double emulsification process, called water-in-oil-in-water (W/O/W) emulsification, is used. The protein solution is first dispersed into a polymer solution dissolved in an organic solvent, and the resulted emulsion is further dispersed into a dilute aqueous solution of another polymer, followed by solvent evaporation or extraction. The major problem associated with the conventional microencapsulation procedures is that the protein molecules may be denaturated by contacting with the organic solvents which are indispensable in the processes. Although surfactants and/or hydrogel solutions are used in the first emulsification to protect the proteins<sup>[4,5]</sup>, they are only effective to certain relatively stable proteins. A microencapsulation process which is free of organic solvents is highly demanded.

For aqueous systems, particles may be formed through various precipitation mechanisms such as salting out<sup>[6]</sup>, acid-base interaction<sup>[7]</sup>, pH assistant precipitation. For these mechanisms, concentrated salts, extreme pH or protein (ionic) cross-link agents are unavoidable. These are all considered chemical hazards to the activity of biological therapeutics.

Two polymer aqueous solutions may be immiscible due to their chain length and structural difference<sup>[8]</sup>. Such polymer aqueous two-phase systems (which are not emulsions but block phases) are practically used for protein purification<sup>[8,9]</sup>. Their aqueous nature and relatively low interfacial tension provide excellent compatibility with soluble proteins in terms of preventing protein conformation change. Protein purification is based on their partition which favors one of the aqueous polymer phase with impurities partitioning into the other. The practice of protein purification is evident that proteins can be distributed into one of the aqueous phases with biological activity intact. This two-phase system readily forms two block phases after mixing. For microencapsulation purpose, however, a stable emulsion must be formed with the two polymer aqueous solutions.

This invention is aimed to address the above-mentioned issues and to develop a new formulation strategy for susceptible therapeutics especially biological agents.

## SUMMARY OF THE INVENTION

The present invention provides a stable aqueous/aqueous emulsion system which is prepared  
5 with a hydrophilic polymer.

This invention also provides the method of preparing a stable aqueous/aqueous emulsion comprising steps of: a) selecting appropriate polymeric materials for  
10 dispersed phase and continuous phase which are immiscible, biocompatible and have biased partition to the active ingredients to be encapsulated; b) selecting appropriate surface modifiers which are charged, non-toxic, and possessing a moderate  
15 interfacial tension between the above two phases; c) developing phase diagram for the above; and d) dispersing the dispersed phase into the continuous phase under an appropriate shear stress.

20 Finally, the invention provides an encapsulation comprising the emulsion system which is prepared with a hydrophilic polymer.

The present invention demonstrates a stable emulsion  
25 system which provides the solution for all the problems raised above by that both the dispersed and the continuous phases are formed from aqueous solutions without concentrated salts, extreme pH, and other chemical hazards.

30 Water soluble proteins, liposomes, live viruses and other therapeutic agents can be microencapsulated on the bases of their partition favoring the dispersed phase, and released or reconstituted upon or prior  
35 to administration with their original morphology and activity preserved.

The emulsion can be dried to fine powder through freeze-drying, spray drying and other methods, and subjected to further treatment: coating, double-microencapsulation, compressing and other procedures  
5 by which a variety of pharmaceutical dosage forms including controlled release systems and targeted delivery systems can be prepared.

For biological agents including live viruses,  
10 encapsulation into the dry form can not only improve their stability and shelf-life, but also avoid expensive cold-chain (-20 °C) in transportation and application. This is important for many developing countries where cold-chain is not available.

15 All the polymer materials used in the invented emulsion system (including dispersed phase, the continuous phase and the surface stabilizing agent) are biocompatible and good for internal use on  
20 humans.

## DETAILED DESCRIPTION OF THE FIGURES

### **Figure 1. Polymer aqueous/aqueous emulsion system.**

Polymer solution A and polymer solution B are  
5 immiscible, thus A can be dispersed into B under a  
shear stress. The third polymer carries charge and  
is fairly immiscible with both A and B at low  
concentration, so that it tends to be rich at the  
interface of A and B, and forms a charged surface.  
10 The charged surface effectively prevents aggregation  
and fusion of the dispersed phase (See *Example 1* in  
the paragraph). Therapeutic agents such as  
proteins, liposomes and viruses are partitioned and  
encapsulated in the dispersed phase and subjected to  
15 lyophilization (See *Examples 2, 3 and 4*).

### **Figure 2. Microscopic image of polymer aqueous/aqueous emulsion.**

20 **Figure 3. Secondary surface modification of polymer aqueous droplets dispersed in a polymer aqueous continuous phase.**

The surface of the dispersed phase in Figure 1 can  
be further modified for functionization (See  
25 *Additional applications* in the paragraph).  
Permeability barrier can be assembled on the surface  
by ionic cross-linking with a degradable polymer  
having opposite charge, or by assembly of a lipid  
bilayer having opposite charge. Release rate of  
30 encapsulated therapeutics can be adjusted by  
selecting the cross-linking polymer in terms of  
chain length and structure of desired degradation  
rate (polyaminoacid or polypeptide for example).  
Targeting moieties (antibodies or ligands) can be  
35 immobilized on the surface through ionic interaction

or hydrophobic interaction (in the case of lipid bilayer assembly).

**Figure 4.        Microspheres prepared by double-**  
**microencapsulation through solid-in-oil-in-water**  
**emulsification**

The powder formed by drying of the aqueous/aqueous emulsion can be further encapsulated into hydrophobic, degradable polymer microspheres. Since methane dichloride, a commonly used solvent in polymer microsphere preparation, dissolves phase B (the continuous phase of the A/A emulsion) but does not interact with phase A (the dispersed phase), the phase B can be removed from the lyophilized powder simply by washing with the solvent. Microspheres which encapsulate the lyophilized powder possess more hydrophobic matrix if the phase B is washed out, but less hydrophobic if the phase B remains. This structural difference can affect degradation rate of the polymer matrix and diffusion rate through the polymer matrix, thus the release profile of encapsulated therapeutics can be adjusted by the content of the phase B remained.

**Figure 5.        Nano-sized preparation using polymer**  
**aqueous/aqueous emulsion.**

Nano-meter-sized crystals and other assemblies formed from two reactants can be prepared using the emulsion system (See reference [14]). Reactant A is usually those which partitioned and encapsulated into the dispersed phase. Reactant B is those which are distributed to both phases. Since A is isolated with limited quantity in each micro-sized droplet, when the assembly process proceeds, the limited accessibility of the reactants ensures a small sized



product. Nano-sized preparation is useful in produce of both therapeutic and diagnostic agents.

5 **Figure 6. Microscopic image of reconstituted AmB/liposomes (of SUV) which were freeze-dried after loading into the polymer emulsion**

Liposomes encapsulated into the polymer emulsion system, followed by lyophilization, are not visible after reconstitution, indicating that their small  
10 unilamellar structure is protected by the polymer emulsion system.

15 **Figure 7. Microscopic image of reconstituted AmB/SUV which were freeze-dried without loading into the polymer emulsion.**

Small liposomes (SUV) after direct lyophilization shows large particles when reconstituted, indicating aggregation and fusion of unprotected liposomes  
20 during the drying process

## Detailed Description of the Invention

The present invention provides a stable aqueous/aqueous emulsion system which is prepared with a hydrophilic polymer. This invention also provides the method of preparing a stable aqueous/aqueous emulsion comprising steps of: a) selecting appropriate polymeric materials for dispersed phase and continuous phase which are immiscible, biocompatible and have biased partition to the active ingredients to be encapsulated; b) selecting appropriate surface modifiers which are charged, non-toxic, and possessing a moderate interfacial tension between the above two phases; c) developing phase diagram for the above; and d) dispersing the dispersed phase into the continuous phase under an appropriate shear stress. This invention further provides the above aqueous/aqueous emulsion system with polymeric surface modifier.

In addition, this invention provides a method for encapsulating protein or peptide comprising the above emulsion system. In an embodiment, the encapsulated protein or peptide is used for sustained release formulations or dry powder formulations.

This invention further provides an encapsulation comprising the emulsion system which is prepared with a hydrophilic polymer. In an embodiment, the encapsulation encapsulates protein, peptide, virus, bacterium, or cell.

This invention also provides a liposome-based drug formulation which comprises the above emulsion system. Still further, the invention provides

viral, bacterial or cell microencapsulation comprising the above emulsion system.

5 This invention also provides the nano-sized preparation comprising the above emulsion system. In an embodiment, the preparation is nano-sized crystallization, nano-sized precipitation or other nano-sized assembly.

10 Finally, this invention provides a diagnosis kit comprising the above emulsion system.

15 The polymer aqueous/aqueous emulsion system described in this invention is fundamentally different from existing emulsions which are prepared by dispersing a hydrophilic phases into a hydrophobic phase or the vice versa. This difference is extended to the so-called water-in-oil-in-water (W/O/W) emulsions. In any of these  
20 conventional emulsions, phase separation occurs between an aqueous phase and a water-immiscible organic phase. In the case of the present invention, phase separation occurs between two aqueous phases.

25 The emulsification process is distinct from those precipitation processes from aqueous solutions which are based on salting out, acid-base interaction, and pH-assisted precipitation. In the invented  
30 emulsification process, such mechanisms which rely on application of concentrated salts and extreme pH are not involved. Instead, phase separation between two polymer aqueous phases is due to positive enthalpy of mixing, ( $\Delta H_M$ ), and reduced entropy of  
35 mixing, ( $\Delta S_M$ ), as polymer chain increased.

This system is also different from polymer aqueous two-phase system which is used in protein purification. In addition to phase separation, the dispersed droplets must be stabilized to prevent fusion. This is accomplished by introducing a third aqueous polymer which is charged and adsorbed at the interface between the two immiscible polymer phases. The surface charges brought by the third polymer effectively prevent aggregation and fusion of the droplets.

In the present invention, applications of the polymer aqueous/aqueous emulsion system in pharmaceutical formulation were examined by microencapsulation of three representative agents: liposomes, proteins and live viruses. The new emulsion system showed encouraging result for each type of the agents, indicating its usefulness in formulating susceptible therapeutic agents.

Liposomes are used for formulating pharmaceutical dosage forms for intravenous injection (IV)<sup>[10]</sup>. For IV administration, the liposomes must be prepared and maintained in the form of small unilamella vesicles (SUV). However, during shelf time, the SUV aggregate and fuse to form large particles. The emulsion system may effectively protect and preserve the SUV structure of liposomes by encapsulating them into the dispersed phase and lyophilizing to dry powder. In our study, the protected SUV structure was reconstituted simply by adding water to the lyophilized powder (see **Figure 6**). As the control, un-encapsulated SUV liposomes were lyophilized and re-hydrated, and found that the original SUV morphology was lost and converted to large multilamella vesicles (see **Figure 7**).

In developing sustained release dosage forms for proteins, preventing denaturation of the therapeutic proteins in the formulation processes is a key issue. Denaturation of proteins in formulation processes is mainly due to contact with organic solvents used and strong interfacial tension between the dispersed and the continuous phases. The present invention offers a formulation environment which is free of these chemical hazards found in conventional microencapsulation processes which are regarded as the cause of protein denaturation. As an example, a conformation-sensitive enzyme, ( $\beta$ -galactosidase, was encapsulated with the polymer aqueous/aqueous emulsification technology and compared its activity with that un-encapsulated (the positive control) and that encapsulated with the conventional W/O method (the negative control). The enzymatic activity of the protein after release was comparable with the positive control, while that formulated with the W/O method showed no activity. The result confirms our hypothesis that protein conformation can be preserved during this new microencapsulation process.

In addition to protection of proteins from contact with organic solvents, the present system offers more useful physical chemical mechanisms in controlling released rate and release profile of encapsulated protein therapeutics. For biodegradable polymeric drug delivery systems, release rate of encapsulated therapeutics depends on the rate of degradation (chemical reaction). However, manipulating the rate of a chemical reaction in vivo is difficult due to the restrictions in changing reaction conditions such as

temperature. For the microspheres made by the solid-in-oil-in-water methods, since the phase B (PEG) can be co-dissolved with the hydrophobic polymer (such as PLGA), the nature of the polymer matrix such as hydrophilicity/hydrophobicity, swellability and the rate of hydrolysis, can be adjusted by the content of the phase B (PEG) in the hydrophobic matrix (see **Figure 4**). Moreover, release of PEG can open channels for dissolution of the phase A (dextran), thus lead to a diffusional mechanism for protein release.

Applications of this new material system have been extended to formulation of live viruses. Live viruses are used in some human vaccines <sup>[11]</sup> and many veterinary vaccines. To maintain the viral activity in infecting cells, cold-chain (-20°C) is required in transportation and application of viral vaccines and other viral products. With the present invention, the viral products may be prepared as dry powder without losing their activity, so that room temperature maintenance will be possible. In a preliminary experiment, cytomegalo viruses (CMV) were encapsulated with the aqueous/aqueous emulsion system, followed by lyophilization. The dried viruses were then reconstituted with buffer and incubated with human foreskin fiber blast cells. The infection activity of the encapsulated viruses was compared with fresh viruses (positive control) and viruses lyophilized without the emulsion system (negative control). Again, an unequivocal result was attained that the encapsulated viruses showed an activity comparable to the positive control while the negative control showed no infection.

## Details of the Invention

### **Example 1: Preparation of Polymer Aqueous/Aqueous Emulsion**

5

#### *Method 1*

Dextran (MW 100,000 to 1,000,000) and sodium alginate (low or medium viscosity) were dissolved in water at the dextran concentration of 10 to 50 w/v% and dextran to alginate ratio of 10:1 to 30:1. This solution is named solution A. An aqueous solution, named solution B, containing polyethylene glycol (PEG, MW 1,000 to 12,000) was prepared with PEG concentration ranging from 10 to 40 w/v%. Solution A was added into solution B at the volume ratio from 1:0.7 to 1:5 under a shear stress (stirring or homogenizing). The particle size distribution, which may be measured using a particle sizer for appropriate range, ranged as a function of the shear stress applied. Stability of the emulsion system was examined by placing it at room temperature for several weeks. Fusion was not observed (See figure 1).

25

#### *Method 2*

Solution A was prepared with Dextran (MW 100,000 to 1,000,000) only at the concentration of 10 to 50 w/v%. Solution B containing PEG and alginate of the above-mentioned molecular weight was prepared. The concentration of PEG ranged from 10 to 40 w/v% and PEG to alginate ratio ranged from 10:1 to 30:1. Solution A was dispersed into solution B with the same procedure as above. Stable emulsions were prepared as method 1.

35

## ***Example 2: Encapsulation of liposomes***

### *Partitioning of Liposomes*

5

Solution A, dextrin 30 w/v% and solution B, PEG 25 w/v% were prepared. Small unilamellar vesicle (SUV) liposomes were prepared by sonication of a phospholipid (DOPC with 2% fluorescent lipid) water suspension (lipid/water = 5 ~ 10 mg/ml). Prior to sonication, the lipid-water suspension was sealed with nitrogen. The sonication was sustained, with an interval for each 2 min. until the milky suspension converted to a transparent liquid phase. The resulted liquid was examined using a microscope, and no visible liposomes were found. The liposome suspension was added into solution A and well mixed, so that solution A became colored (yellow). Then solution A with liposome suspension was added to same solution B of the same volume, followed by stirring for 10 min. After the emulsified solution (containing A and B) was allowed to settle for 10 min, the cloudy emulsion became two clear block phases. The dextran phase, which at the bottom, was yellow, while the PEG phase was colorless. This is evident that the liposomes are mainly distributed in the dextran phase.

### *Encapsulation and lyophilization of liposomes*

30

Solution A and B, same as in Example 1, were prepared. Small liposomes (SUV) were prepared as above without fluorescent lipids. The liposomes were first dispersed in solution A, followed by further emulsification with solution B. The resulted emulsion was allowed to settle over night,

35



and no precipitation was observed. The sample was then frozen and subjected to a lyophilizer with vacuum better than  $10^{-2}$  torr for over night.

5 The resulted dry powder was reconstituted by readily dissolving in water. The solution was clear and no visible liposomes were found under a microscope (See **Figure 6**). For comparison, the same liposomes suspension was lyophilized directly without  
10 dispersing into the polymer solutions. Adding water to the directly lyophilized powder resulted in a milky suspension, and large (visible) liposomes were identified under a microscope (See **Figure 7**). This experiment indicates that aqueous emulsion system  
15 can effectively protect the SUV structure from collapse during lyophilization process. The liposome (SUV) suspension was also dispersed into a dextran solution (solution A in Example 3), followed by lyophilization. Instead of fine powder, a hard block  
20 was resulted from this procedure. Dissolving the block sample required vortex for approximately 5 min.

**Example 3: Activity of  $\beta$ -galactosidase after**  
25 **microencapsulation**

*Encapsulation of  $\beta$ -galactosidase*

Beta-galactosidase (1000 unit/ml), 2  $\mu$ l, was added  
30 into 0.25 ml solution A (same as in example 1), and mixed with pipetting. The resulted solution was dispersed slowly into solution B (example 1) under stirring at room temperature. The volume ratio of solution A and B was 1:1.

35

Encapsulation efficiency of the protein in the dispersed phase (the dextran phase) was examined by enzymatic activity of the protein in hydrolysis of 0-nitrophenyl-beta-D-galactopyranoside. Prior to enzymatic activity test, the dispersed phase was separated from the continuous phase by centrifugation at 500 G for 2 minutes.

A solution containing 30 mM Mg, 50 mM sodium phosphate, and 10 mM o-nitrophenyl-beta-D-galactopyranoside (the substrate) was prepared for assay of the enzymatic activity. This reactant solution, 0.3 ml, was mixed with each 0.5 ml of the dextran and PEG phases separated as above, respectively. The two assay samples were then incubated at 37 °C for 30 minutes, followed by addition of 0.5 ml 1 M Na<sub>2</sub>CO<sub>3</sub> into each of them to terminate the reaction. Since the product of the reaction shows yellow color, the samples were subjected to a photometer and absorption at 420 nm was recorded. The absorbance was 0.82 and 0.08 for the dextran phase and the PEG phase, respectively. This result indicates that encapsulation efficiency is approximately 90%.

#### *Protection of protein activity*

To examine the compatibility of this emulsification process with conformational sensitive proteins, the enzymatic activity of encapsulated  $\beta$ -galactosidase was compared with two references. One was a positive control that the  $\beta$ -galactosidase solution was added into the reactant solution directly. The other was a negative control that 2  $\mu$ l of the enzyme solution was added to 0.25 ml solution A (prepared

as in example 1), followed by dispersing into a mineral oil. The protein in the W/O emulsion was recovered by washing the oil away with acetone and re-dissolving the pellet. After adding the reactant solution and incubation as above, the absorbance recorded was 0.81 and 0.00 for the positive and the negative controls, respectively. The result is unequivocal that the enzymatic activity of  $\beta$ -galactosidase was preserved during the microencapsulation process with the aqueous/aqueous emulsification but destroyed with the conventional W/O processes.

#### ***Example 4: Dry formulation of live viruses***

##### *Microencapsulation of cytomegalo viruses*

Microencapsulation of cytomegalo viruses (CMV) was carried out under a procedure similar to that for liposome in example 2. Solution A containing 25 w/v% dextran 500T, 2 w/v% sodium alginate (medium viscosity), 50mM Tris and 100 mM sodium chloride, and solution B contains 25 w/v% PEG 8000 were prepared for microencapsulation. CMV suspension, 5  $\mu$ l in arbitrary unit was mixed with 0.4 ml of solution A, and then dispersed into 0.3 ml of solution B under magnetic stirring. While the initial volume of solution A was larger than that of solution B, the former still formed the dispersed phase. The emulsified samples were subjected to lyophilization as for liposomes in example 2 to form dry powders.

*Infection activity*

The assay viral activity in infecting cells, four samples were prepared:

5

1) CMV in TN buffer (50 mM Tris, 100 mM NaCl),  
without lyophilization; (+ control)

2) CMV in TN buffer, lyophilized; (- control)

10

3) CMV microencapsulated but without  
lyophilization;

4) CMV microencapsulated and lyophilized;

15

After lyophilization, sample 2 and 4 were placed at room temperature for a day, and reconstituted by adding more TN buffer which dissolved the polymer powder. Then the four samples were incubated with human foreskin fiber blast cells. Each of the sample was diluted to six concentration (ten times different between two adjacent concentrations) and added to six dishes of the cells, respectively. The infections were monitored by formation of patches of the linear cells on the dishes. After incubation for a week, microencapsulated viruses (sample 3 and 4) showed comparable activity in infecting cells, but the activity was lower than that of the positive control (sample 1). The negative control (sample 2), however, showed no infection on any of the six dishes. The result is evident that the microencapsulation approach can protect viruses during lyophilization and lead to an active dry powder viral formulation.

35

### ***Additional applications***

#### *Controlled release and targeted delivery of therapeutics*

5

The charged surface of the dispersed droplets can be further modified through ions cross-linking with a polymer of opposite charge (Figure 2). Such charged polymers which are degradable and biocompatible are available (polypeptides, polyaminoacids [5], and chitosan, for example). By selecting the cross-linking agents (in terms of degradation rate and dissolution rate), release rate of encapsulated therapeutic agents can be adjusted.

15

The surface charge can be used to immobilize targeting moieties on the surface for cell-targeted delivery of therapeutics. Ligands or antibodies can be co-polymerized with a charged polyaminoacid [5], and immobilized on the particle surface (See Figure 2).

20

The surface charge can also be used to assemble a phospholipid bilayer which enclosed the particle<sup>[12, 13]</sup>. A supported phospholipid bilayer provides bio-function and cell surface environment to an artificial surface<sup>[14]</sup> which allows functional membrane proteins be immobilized and reconstituted on the surface.

25  
30

#### *Nano-sized preparation*

Nano-meter-sized crystals and other solid particles are actively used in therapeutic (nano-cochleates [14]) and diagnostic (colloidal gold) agents. With the aqueous/aqueous emulsion system, one of

35

reactants may be isolated into the micro-droplets waiting for the other reactant, which may be ions or other precipitation/crystallization agents, to diffuse into the droplet and initiate the reaction.

- 5 Because one of the reactant is isolated in the micro-sized droplets, the growth of the crystal or other assembly is limited due to the limited accessibility between the reactant molecules.

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
22

## REFERENCES

1. R. Langer and J. Folkman, "Polymers for the  
5 sustained release of proteins and other  
macromolecules"; *Nature*, 263, 797-800 (1976)
2. S. Cohen and H. Berstein, *Microparticulate  
systems for delivery of Proteins and vaccines*;  
10 Marcel Dekker, New York, 1996
3. R. Bodmeier, H. Chen, P. Tyle and P. Jarosz,  
"Pseudo phedrine Hcl microspheres formulated  
into an oral suspension dosage form"; *J.*  
15 *controlled Rel.*, **15**, 65-77 (1991)
4. M.J. Alonso, R.K. Gupta, C. Min, G.R. Siber and  
R. Langer, "Biodegradable microspheres as  
controlled release tetanus toxoid delivery  
20 systems"; *Vaccine*, **12**, 299 (1994)
5. T. Jin and K. Solkoll, un-published data
6. B.E. Conway, *Ionic hydration in chemistry and*  
25 *biophysics*; Elsevier, Amsterdam, 1981
7. A.R. Cross and C. Anthony, *Biochem. J.*, 192,  
421 (1980)
- 30 8. B.Z. Zaolavsky, *Aqueous Two-phase partitioning*;  
Marcel Dekker, New York, 1994
9. P.A. Albertsson, *Partition of cell particles  
and macromolecules* 3rd ed. Wiley, New York,  
35 1996

10. AmBisomeR, (Developed by NeXstar),  
www.gileed.com
- 5 11. N.T. Parkin, P. Chiu, and R. Coelinph,  
"Genetically-engineered live attenuated  
influenza A virus vaccine candidates"; *J.*  
*Viro.*, **74**, 2772-2778 (1997)
- 10 12. E. Sakmann, *Science*, 271, 43-48, (1996)
13. R.J. Lee, L. Huang, *Artificial Self-Assembling  
Systems for Gene Transfer*, P.L. Felgner et al.  
ed., American Chemical Society, Washington DC,  
1996
- 15 14. L. Zarif, I. Segarra, T. Jin, D. Hyra, R.J.  
Mannino; "Amphotericin B cochleates as a novel  
oral delivery system for the treatment of  
fungal infections," *26th International  
Symposium on Controlled Release of Bioactive  
Materials*; Boston, USA, 1999
- 20